CAMP EMERGENCY MEDICATION PLAN

Scout'sName:		_ Date of Birth:	Pack or Troop #:
Cam	p Hinds Cam	p Telephone & Fax:	207-655-4878
TO BE COMPLETED BY PARENT OR GUARDIAN:			
I authorize the exchange of medical	information about my child	's asthma between the	physician's office and camp nurse.
Parent or Guardian signature:			
Parent or Guardian tel.# home:	work:_		_ cell phone:
Physician/Healthcare Provider Name:		Parent concerns:	
My child may carry and use his/her:	inhaled asthma medicine	□ Yes □ No	Epi-Pen □ Yes □ No □ N/A
TO BE COMPLETED BY CAMPER		•	
Provider name:			Fax#
	□ NO changes	from previous pla	an
Peak Flow:			
Child's predicted, or personal bes			
Child's Green Zone:	Yellow Zone:		Red Zone: below
Medications:			
	ications.		
Preventive (Controller) Medi	ications:		
Quick Relief Medications: (c.	heck the appropriate quick re	elief med, circle device, l	ist dose/ frequency):
☐ Albuterol (Proventil, Ventolin)	☐ Pirbuterol (Maxair)	☐ Other:	·
Inhaler with spacer OR nebulize	er Dose/Frequency:		
Allergies /Triggers for asthn	na: □ None known		
□ Avoid animals	IIa. Li None anomi		
☐ Other triggers to avoid:			
Exercise Pretreatment Instru			
			or
			child's peak flow is below
Asthma Exacerbation Treatr	ment Instructions:		
> YELLOW ZONE: If child is cou		throath and/or neak f	lawia in Vallaw Zana
		· · · · · · · · · · · · · · · · · · ·	be repeated in 10 minutes if doesn't
recover to Green Zone. Notif	fy parents of exacerbation.	Julizer deadrionic, may	be repeated in 10 minutes in accent
☐ Other:			
RED ZONE: If child is in respi	iratorv distress, and/or pea	k flow is in Red Zone:	
☐ Give 4 puffs quick relief inhaler	r (or nebulizer treatment), and	d call parent and Healtho	
Call 911 if child does not im	prove quickly or parents/He		
Other:			
Special Instructions:			
Maine law now permits campers to	o carry and use inhaled me	dications and Epi-pen	after demonstrating appropriate use of
Inhalers and or Epi-Pen to camp n			= - · · · · · · · · · · · · · · · · · ·
This camper has the knowledge	•		
This camper is not able to carry	-		
 Please contact Healthcare Provi of pre-exercise treatment) 	ider and parent ii camper is u	sing quick relief medicii	nes more than 2 times a week (i.e. in excess
Other:			
Healtho	care Provider signature	D	ate
TO BE COMPLETED BY CAMP NURSE: This camper demonstrates knowledge and skill to carry and use:			
Inhaler medications	□ NO _		
Epi-Pen ☐ YES	□ NO □ N/A	Camp Nurse Sign	ature Date

CAMP EMERGENCY MEDICATION PLAN

ASTHMA PLAN INSTRUCTIONS

Every camper with asthma should have a current

Camp Emergency Medication Plan completed and signed by their physician (or other health care provider) and kept on file in the camp nurse's office. The form must also be signed by a parent/guardian. The plan should be updated each year or when there are major changes to the plan (such as in medication type or dose). The physician's office is encouraged to fax the plan to the camper's camp nurse.

The camp plan is intended to strengthen the partnership of families, healthcare providers and the camp. It is based on the NHLBI Guidelines for Asthma Management. (For more information contact the camp nurse or www.nhlbi/nih/gov).

CARRYING AND ADMINISTERING QUICK RELIEF INHALERS and/or Epi-Pen:

- Most campers are capable of carrying and using their quick relief inhaler by themselves. The camper, camper's parents, camp nurse and healthcare provider should make this decision. The camp nurse must also evaluate technique for effective use.
- The appropriate boxes must be checked by the parent, provider and camp nurse to indicate the camper's ability to carry and self-administer these medications.

USE OF QUICK RELIEF MEDICATIONS MORE THAN TWICE WEEKLY:

> This indicates poor control of asthma, and providers should be notified by the camp nurse or designated staff.

PEAK FLOW ZONES (based on camper's personal or predicted best):

Green zone: Peak flow 80-100%

- ➤ Symptoms and/or use of quick relief medication < 2 times a week
- > Use daily controller medication at home
- > Full participation in physical education and sports

Yellow zone: Peak flow 50-80%

- ➤ Has symptoms or needs quick relief medication >2 times a week
- > Needs quick relief medication and further observation by camp nurse; notify parents
- > Attend physical education but restrict strenuous aerobic activity

Red zone: Peak flow <50%

- Symptoms may include shortness of breath, retractions, difficulty talking or walking; quick relief medication not effective
- > Requires immediate action, close monitoring and notification of parent and healthcare provider